



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

Highlights of Novel Vaccination Strategies in Allergen Immunotherapy

Komlósi, Zsolt István ; Kovács, Nóra ; Sokolowska, Milena ; van de Veen, Willem ; Akdis, Mübeccel ; Akdis, Cezmi A

Abstract: Increasing safety while maintaining or even augmenting efficiency are the main goals of research for novel vaccine development and improvement of treatment schemes in allergen immunotherapy (AIT).^{1, 2, 3, 4} In spite of encouraging positive experiences,⁵ AIT still faces several problems related to worldwide standardization, its limited efficacy, potential severe side effects, low patient adherence, high costs, and long duration (3–5 years) of treatment. There have been many different approaches to improve the standardization, efficacy, and safety of AIT in the past, which have been continuously pursued during the past years.^{6, 7, 8} Currently, efforts on developing novel vaccines of AIT continue for the treatment of asthma,⁹ allergic rhinoconjunctivitis,¹⁰ bee venom allergy,¹¹ and food allergy.¹² Similar efforts are being undertaken for autoimmune diseases. Studies to offer prophylactic usage are also being performed.¹³ Regulatory issues represent an important aspect in this context¹⁴ and global and national strategies are being continuously developed.^{15,16} It is essential to develop strong collaborations with the regulatory authorities for extensive preventive usage of AIT vaccines and usage of pollen chambers for outcome evaluations to avoid slowing down of vaccine developments. Here, the authors summarize some current investigations aiming at further improvement of AIT vaccines (Table 1).

DOI: <https://doi.org/10.1016/j.iac.2019.09.010>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-180549>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Komlósi, Zsolt István; Kovács, Nóra; Sokolowska, Milena; van de Veen, Willem; Akdis, Mübeccel; Akdis, Cezmi A (2020). Highlights of Novel Vaccination Strategies in Allergen Immunotherapy. *Immunology and Allergy Clinics of North America*, 40(1):15-24.

DOI: <https://doi.org/10.1016/j.iac.2019.09.010>

Highlights of Novel Vaccination Strategies in Allergen Immunotherapy



Zsolt István Komlósi, MD, PhD^{a,*}, Nóra Kovács, MD^{a,b},
Milena Sokolowska, MD, PhD^{c,d}, Willem van de Veen, PhD^{c,d},
Mübeccel Akdis, MD, PhD^{c,d}, Cezmi A. Akdis, MD^{c,d}

KEYWORDS

- Allergy • Allergen immunotherapy • AIT • Adjuvants • Allergoids
- Recombinant allergen peptides • Active immunization • Passive immunization

KEY POINTS

- The current aim is to improve allergen immunotherapy (AIT) to shorten duration of treatment, enhance efficiency, reduce side effects, and, ultimately, significantly increase the utilization of this curative treatment of allergy with high patient adherence and compliance.
- To increase the efficacy of AIT, allergens have been modified to facilitate their uptake and presentation, or have been coupled to innate immunostimulatory substances. New adjuvants have also been introduced recently.
- Hypoallergenic molecules (eg, allergoids and recombinant allergen peptides) have been developed to improve the safety profile of the vaccines.
- Administration of recombinant IgG₄ antibodies is a new, quick, passive immunization AIT strategy with remarkable efficiency.
- Recent mouse experiments suggest that it may be possible to develop AIT even for pre-natal allergy prevention.

INTRODUCTION

Increasing safety while maintaining or even augmenting efficiency are the main goals of research for novel vaccine development and improvement of treatment schemes in allergen immunotherapy (AIT).^{1–4} In spite of encouraging positive experiences,⁵ AIT still faces several problems related to worldwide standardization, its limited efficacy,

^a Department of Genetics, Cell- and Immunobiology, Semmelweis University, Nagyvárad Sqr. 4, Budapest 1089, Hungary; ^b Lung Health Hospital, Munkácsy Mihály Str. 70, Törökbálint 2045, Hungary; ^c Swiss Institute of Allergy and Asthma Research (SIAF), Hermann-Burchard Strasse 9, Davos Wolfgang CH7265, Switzerland; ^d Christine Kühne – Center for Allergy Research and Education, Hermann-Burchard Strasse 1, Davos Wolfgang CH7265, Switzerland

* Corresponding author.

E-mail addresses: komlosi.zsolt@med.semmelweis-univ.hu; drkomlo@yahoo.com

potential severe side effects, low patient adherence, high costs, and long duration (3–5 years) of treatment. There have been many different approaches to improve the standardization, efficacy, and safety of AIT in the past, which have been continuously pursued during the past years.^{6–8} Currently, efforts on developing novel vaccines of AIT continue for the treatment of asthma,⁹ allergic rhinoconjunctivitis,¹⁰ bee venom allergy,¹¹ and food allergy.¹² Similar efforts are being undertaken for autoimmune diseases. Studies to offer prophylactic usage are also being performed.¹³ Regulatory issues represent an important aspect in this context¹⁴ and global and national strategies are being continuously developed.^{15,16} It is essential to develop strong collaborations with the regulatory authorities for extensive preventive usage of AIT vaccines and usage of pollen chambers for outcome evaluations to avoid slowing down of vaccine developments. Here, the authors summarize some current investigations aiming at further improvement of AIT vaccines (Table 1).

NOVEL ALLERGEN FORMULATIONS AND ADJUVANTS

Adjuvants are add-on constituents of the vaccines, ideally increase the immunogenicity of the allergens to induce a quicker, stronger, and longer-lasting immune response; meanwhile also modulate the immune response to enhance allergen-specific T regulatory cell differentiation, immunoglobulin (Ig)G₄ antibody production, and, importantly, avoid T_H2-type response as well as anaphylaxis.

Coupling of allergens to innate immunostimulatory substances is primarily aimed at improving vaccine efficiency. These strategies include CpG oligonucleotide-conjugated allergens,¹⁷ allergens coupled to viruslike particles,^{18,19} carbohydrate-based particles,²⁰ and the use of monophosphoryl lipid A formulated with allergoid.²¹ Encapsulation of allergens in nanoparticles enables us to design the immunomodulatory and depot effect of a complex vaccine very precisely.^{22,23} Nanoparticles can have an adjuvant effect themselves (eg, viruslike particles), or various adjuvant molecules can be conjugated to or packed in the particles (eg, *Vibrio cholerae* neuraminidase on synthetic microparticles), and the physical-chemical characteristics of the particles determine the kinetics of the antigen release (eg, liposomes). Liposomes and microcrystalline tyrosine are promising new alternatives of the widely used aluminum hydroxide (alum) adjuvant.

Sublingual administration of an allergen together with α -galactosylceramide-liposomes increased the therapeutic performance of the vaccine in mouse model of allergic rhinitis.²⁴ High-affinity glycan ligand of the inhibitory receptor CD33 and the allergen, packed in liposomal nanoparticles, was able to suppress immunoglobulin (Ig)E-mediated activation of mast cells and, thus, was shown to be efficient in preventing anaphylaxis in transgenic mice bearing human CD33-expressing mast cells.²⁵ Such approaches may help to increase the safety of AIT.

Microcrystalline tyrosine, an alternative depo adjuvant composed of nonessential amino acid L-tyrosine, is a second-generation adjuvant currently used for AIT in humans (eg, Pollinex, Acarovac Plus).²⁶ It is proven to be safe for human use,²⁷ and efficiently induces IgG₄ production and reduces symptoms in house dust mite (HDM) subcutaneous immunotherapy (SCIT).²⁶

Microcrystalline tyrosine-adsorbed grass pollen allergoid (Pollinex quattro) enhanced with monophosphoryl lipid A, a second-generation immunomodulatory adjuvant, had a beneficial long-term effect as well.²⁸

Recombinant allergen peptides, selected to have advantageous safety profile are generally less immunogenic. Therefore, various carrier systems, typically of microbial origin, have been developed.²⁹ Promising new approaches include *Vibrio cholerae*

Table 1 Novel AIT vaccine developmental strategies currently being pursued	
Type of the Vaccine/Approach	References
Coupling of allergens to innate immunostimulatory substances	
CpG oligonucleotide-conjugated allergens	17
Allergens coupled to viruslike particles	18,19,32,33
Carbohydrate-based particles	20
<i>Vibrio cholerae</i> neuraminidase on allergen-loaded synthetic microparticles	29
Fusion of nonallergenic peptides and PreS protein from hepatitis B virus	34,35
Fusion of nonallergenic peptides and tetanus toxoid	36
Novel adjuvants	
Monophosphoryl lipid A formulated with allergoid	21
Microcrystalline tyrosine	26,27
Proteoliposome adjuvant from <i>Neisseria meningitidis</i>	30
α -Galactosylceramide containing liposomes	24
Encapsulation of allergens in micro-/nanoparticles	
Carbohydrate-based particles	20
α -Galactosylceramide containing liposomes	24
CD33 ligand and the allergen packed in liposomal nanoparticles	25
<i>V. cholerae</i> neuraminidase on allergen-loaded synthetic microparticles	29
Allergoids and recombinant allergens	
Monophosphoryl lipid A formulated with allergoid	21
Allergen extracts modified with glutaraldehyde	40
Fusion of nonallergenic peptides and PreS protein from hepatitis B virus	34,35
Fusion of nonallergenic peptides and tetanus toxoid	36
Recombinant T-cell epitope peptides	43
Allergen fragments	44
Recombinant allergen hybrids	45
Conjugation of allergoids to nonoxidized mannan	51,52
Fusion of allergens with other immune response modifiers	
CD33 ligand and the allergen packed in liposomal nanoparticles	25
MAT-vaccine	37,38
Conjugation of allergoids to nonoxidized mannan	51,52
Novel routes of administration	
Intralymphatic AIT	37,38,60
Epicutaneous AIT	59
Combination of AIT with biologicals	
Administration of anti-IgE	55–58
Novel administration protocols	
Passive immunization	69
Prenatal preventive vaccination	68,70

Considering that certain approaches may include different strategies, some of the vaccines are mentioned in multiple categories.

Abbreviations: AIT, allergen immunotherapy; Ig, immunoglobulin; MAT, modular antigen translocation.

neuraminidase, as an immunomodulatory agent, on allergen-loaded synthetic micro-particles³⁰; proteoliposome from *Neisseria meningitidis* as an adjuvant for HDM allergy vaccine³¹; and utilization of viruslike particles to enhance uptake of the allergen by antigen-presenting cells.³² A single dose of allergen Fel d 1-linked virus-like particles is able to prevent type I hypersensitivity response in mice.^{33,34} Fusion proteins consisting of nonallergenic peptides from the 4 major timothy grass pollen allergens and the PreS protein from hepatitis B virus as a carrier efficiently reduced T-cell proliferation and proinflammatory cytokine release, while increased blocking IgG activity.³⁵ Engineered recombinant fusion proteins consisting of nonallergenic peptides of the allergens and microbial components, like hepatitis B PreS protein³⁶ or partial fragment of a tetanus toxoid molecule,³⁷ seems to be safe and efficient in model systems.

In a novel approach, the major cat allergen (Fel d 1) was modified in a way that its uptake and presentation by professional antigen-presenting cells was highly improved. Human immunodeficiency virus transactivator of transcription (TAT)-derived membrane translocation domain was used to enhance entry to cells, and a truncated peptide of the invariant chain was used to increase antigen presentation.^{38,39} This modular antigen translocation (MAT)-Fel d 1 vaccine is efficiently internalized and potently presented to T cells by antigen-presenting cells, which stimulated T-cell responses in 100 times lower doses compared with native allergens. In a double-blind, placebo-controlled clinical trial, the MAT-Fel d 1 vaccine was administered in 3 increasing doses into inguinal lymph nodes at 4-week intervals. In addition to a good safety profile, only 3 doses of MAT-Fel d 1 intra-lymph node vaccination induced clinical tolerance to nasal cat allergen challenge in parallel with T-cell tolerance and increased serum IgG₄ and interleukin (IL)-10 levels.^{38,39} Currently this approach is being pursued for veterinary vaccine development.

ALLERGOIDS, RECOMBINANT ALLERGENS

Targeting T cells to induce T-cell tolerance and bypassing IgE binding to avoid IgE-mediated side effects is an essential approach in AIT.⁴⁰ To improve AIT safety, the primary aim is to use hypoallergenic molecules to prevent severe complications such as anaphylaxis. Currently, there are many modified allergen preparations (allergoids) in the market with decreased IgE binding activity. Chemical modification of allergen extracts (eg, with glutaraldehyde⁴¹) aims to prevent IgE cross-link with conformational epitopes, while linear epitopes are preserved. To accomplish this aim, several successful new approaches have been developed,^{42,43} including mixture of T-cell epitope peptides,⁴⁴ allergen fragments,⁴⁵ fusion proteins,^{36,37} and recombinant allergen hybrids.⁴⁶ The prototype of this approach is peptide immunotherapy that uses linear T-cell epitope peptides, and has shown promising results,^{47,48} although it still faces challenges in phase III clinical trials.^{49–51} All of these allergoid preparations enable the administration of the allergens in higher doses to efficiently induce T-cell tolerance without the risk of anaphylaxis.⁴⁰ Conjugation of allergoids to nonoxidized mannan facilitates their uptake by antigen-presenting cells.^{52,53}

Some of the clinically significant allergens, however, may be poorly represented in allergen extracts. Development of molecular vaccines, based on recombinant allergen components, helps to overcome the limitations of the vaccines based on natural allergen extracts.⁵⁴ Identification of immunodominant molecular components of the allergens is an important initial step of the development of such AIT approaches.⁵⁵

ADD-ON THERAPIES AND NEW PROTOCOLS

Combination of conventional or novel methods of AIT with biological immune response modifiers is a promising new therapeutic approach. Anti-IgE combined with AIT has been evaluated in several studies. Significant decrease in the risk of anaphylaxis and improved rescue medication scores were observed during rush immunotherapy with a quicker dose increment and reach of the maintenance dose.^{56,57} Accumulating evidence suggests that anti-IgE treatment is a valuable add-on to AIT for airway allergies, especially to prevent adverse events in the build-up phase and improve safety profile in cases in which the designated maintenance dose is not tolerated by the patient.^{58,59} Other possible combinations include anti-IL-4 or anti-IL-13 as well as their receptor antagonists.

Different routes of vaccine administration also have been proposed to improve efficacy and safety. Intralymphatic vaccination and epicutaneous vaccination are recently pursued novel strategies. Both showed similar efficacy to SCIT in grass pollen allergy, however, that has been reached with fewer injections and lower total allergen doses.⁶⁰ Intralymphatic administration induces T-cell tolerance or strong T_H1 responses depending on the type of vaccine used.^{39,61}

Meta-analysis of the double-blind, placebo-controlled trials has shown that sublingual immunotherapy (SLIT) is safe and clinically efficacious with a treatment benefit approximately half of that achieved with SCIT.⁶² Disease-modifying effects of SLIT have been confirmed in large-scale randomized, double-blind, placebo-controlled trials also in children^{63–68}; and even its preventive administration has been proposed.⁶⁹

In an exciting, current study, cat-allergic patients were treated with a single dose of 2, fully human, recombinant IgG₄-blocking antibodies specific for Fel d 1, the major cat allergen. This treatment resulted in a rapid and sustained reduction in clinical symptoms after nasal allergen provocation test, suggesting a new, quick, passive AIT strategy for allergies.⁷⁰ Treatment-emergent adverse events were closely monitored during the study, and there was neither any difference in the frequency of the adverse events between antibody-treated and placebo group, nor did any severe adverse event occur in relation to the IgG₄ antibody treatment.⁷⁰

In a recent mouse experiment, the administration of anti-IgE antibody during pregnancy caused long-term IgE-class-specific immunosuppression in the offspring that was able to prevent allergic sensitization. These results suggest that it may be possible to develop a vaccination strategy aiming prenatal allergy prevention.⁷¹

Reduced side-effects

- Chemically modified allergen
- Recombinant allergen peptides
- Conjugation to nonoxidized mannan
- CD33 ligand

Improved adjuvants

- Microcrystalline tyrosine

New routes and add-ons

- Intralymphatic administration
- Epicutaneous administration
- Single dose recombinant IgG₄
- Add-on anti-IgE

Ideal AIT vaccine

Encapsulation

- Liposomes
- Carbohydrate-based particles
- Virus-like particles

Addressing constructs

- MAT vaccine

Conjugation to microbial products

- Monophosphoryl lipid A
- GpG oligonucleotide
- Hepatitis B PreS protein
- α -galactosylceramide
- *Vibrio cholerae* neuraminidase
- *Neisseria meningitidis* proteoliposome
- Tetanus toxoid fragment

Fig. 1. Strategies for development of a more safe, efficient, and reliable vaccine for AIT.

SUMMARY

Intensive efforts have been invested to improve AIT vaccines by making them more efficient, convenient, and safe^{2,16} (**Fig. 1**). Still, fewer than 10% of allergic patients choose to receive AIT.⁷² Besides the fear from potential side effects and poor adherence due to long duration of treatment,⁷³ the possible cause of weak overall, population-wide efficiency of AIT is that not all the potential influencing factors were considered during the development of AIT strategies, because of lack of knowledge. Identification of good biomarkers predicting beneficial vaccine immune response is still an unmet clinical need; however, it is under intensive investigation.⁷⁴ Better understanding of the immune response to vaccines will hopefully enable us to develop more efficient AIT strategies specifically tailored to allergic disease endotypes.⁷⁵

ACKNOWLEDGMENTS

The laboratory of Z.I. Komlósi is supported by Hungarian Pediatric Oncology Network. The laboratories of C.A. Akdis and M. Akdis are supported by the Swiss National Science Foundation.

REFERENCES

1. Casale TB, Stokes JR. Immunotherapy: what lies beyond. *J Allergy Clin Immunol* 2014;133(3):612–9 [quiz: 620].
2. Agache I. EAACI guidelines on allergen immunotherapy—out with the old and in with the new. *Allergy* 2018;73(4):737–8.
3. Pfaar O, Bonini S, Cardona V, et al. Perspectives in allergen immunotherapy: 2017 and beyond. *Allergy* 2018;73(Suppl 104):5–23.
4. Pfaar O, Lou H, Zhang Y, et al. Recent developments and highlights in allergen immunotherapy. *Allergy* 2018;73(12):2274–89.
5. Zielen S, Devillier P, Heinrich J, et al. Sublingual immunotherapy provides long-term relief in allergic rhinitis and reduces the risk of asthma: a retrospective, real-world database analysis. *Allergy* 2018;73(1):165–77.
6. Passalacqua G, Baena-Cagnani CE, Bousquet J, et al. Grading local side effects of sublingual immunotherapy for respiratory allergy: speaking the same language. *J Allergy Clin Immunol* 2013;132(1):93–8.
7. Jones SM, Burks AW, Dupont C. State of the art on food allergen immunotherapy: oral, sublingual, and epicutaneous. *J Allergy Clin Immunol* 2014;133(2):318–23.
8. Vickery BP, Scurlock AM, Kulis M, et al. Sustained unresponsiveness to peanut in subjects who have completed peanut oral immunotherapy. *J Allergy Clin Immunol* 2014;133(2):468–75.
9. Wahn U, Bachert C, Heinrich J, et al. Real-world benefits of allergen immunotherapy for birch pollen-associated allergic rhinitis and asthma. *Allergy* 2019;74(3):594–604.
10. Roberts G, Pfaar O, Akdis CA, et al. EAACI guidelines on allergen immunotherapy: allergic rhinoconjunctivitis. *Allergy* 2018;73(4):765–98.
11. Sturm GJ, Varga EM, Roberts G, et al. EAACI guidelines on allergen immunotherapy: hymenoptera venom allergy. *Allergy* 2018;73(4):744–64.
12. Pajno GB, Fernandez-Rivas M, Arasi S, et al. EAACI Guidelines on allergen immunotherapy: IgE-mediated food allergy. *Allergy* 2018;73(4):799–815.
13. Holt PG, Sly PD, Sampson HA, et al. Prophylactic use of sublingual allergen immunotherapy in high-risk children: a pilot study. *J Allergy Clin Immunol* 2013;132(4):991–3.e1.

14. Bonertz A, Roberts GC, Hoefnagel M, et al. Challenges in the implementation of EAACI guidelines on allergen immunotherapy: a global perspective on the regulation of allergen products. *Allergy* 2018;73(1):64–76.
15. Larenas-Linnemann DES, Antolin-Amerigo D, Parisi C, et al. National clinical practice guidelines for allergen immunotherapy: an international assessment applying AGREE-II. *Allergy* 2018;73(3):664–72.
16. Muraro A, Roberts G, Halken S, et al. EAACI guidelines on allergen immunotherapy: executive statement. *Allergy* 2018;73(4):739–43.
17. Creticos PS, Schroeder JT, Hamilton RG, et al. Immunotherapy with a ragweed-toll-like receptor 9 agonist vaccine for allergic rhinitis. *N Engl J Med* 2006; 355(14):1445–55.
18. Kundig TM, Senti G, Schnetzler G, et al. Der p 1 peptide on virus-like particles is safe and highly immunogenic in healthy adults. *J Allergy Clin Immunol* 2006; 117(6):1470–6.
19. Engeroff P, Caviezel F, Storni F, et al. Allergens displayed on virus-like particles are highly immunogenic but fail to activate human mast cells. *Allergy* 2018; 73(2):341–9.
20. Thunberg S, Neimert-Andersson T, Cheng Q, et al. Prolonged antigen-exposure with carbohydrate particle based vaccination prevents allergic immune responses in sensitized mice. *Allergy* 2009;64(6):919–26.
21. Rosewich M, Schulze J, Eickmeier O, et al. Tolerance induction after specific immunotherapy with pollen allergoids adjuvanted by monophosphoryl lipid A in children. *Clin Exp Immunol* 2010;160(3):403–10.
22. Di Felice G, Colombo P. Nanoparticle-allergen complexes for allergen immunotherapy. *Int J Nanomedicine* 2017;12:4493–504.
23. Pohlit H, Bellinghausen I, Frey H, et al. Recent advances in the use of nanoparticles for allergen-specific immunotherapy. *Allergy* 2017;72(10):1461–74.
24. Suzuki S, Sakurai D, Sakurai T, et al. Sublingual administration of liposomes enclosing alpha-galactosylceramide as an effective adjuvant of allergen immunotherapy in a murine model of allergic rhinitis. *Allergol Int* 2019;68(3):352–62.
25. Duan S, Koziol-White CJ, Jester WF Jr, et al. CD33 recruitment inhibits IgE-mediated anaphylaxis and desensitizes mast cells to allergen. *J Clin Invest* 2019;129(3): 1387–401.
26. Roger A, Depreux N, Jurgens Y, et al. A novel microcrystalline tyrosine-adsorbed, mite-allergoid subcutaneous immunotherapy: 1-year follow-up report. *Immunotherapy* 2016;8(10):1169–74.
27. Baldrick P, Richardson D, Wheeler AW. Review of L-tyrosine confirming its safe human use as an adjuvant. *J Appl Toxicol* 2002;22(5):333–44.
28. Zielen S, Gabrielpillai J, Herrmann E, et al. Long-term effect of monophosphoryl lipid A adjuvanted specific immunotherapy in patients with grass pollen allergy. *Immunotherapy* 2018;10(7):529–36.
29. Zahirovic A, Lunder M. Microbial delivery vehicles for allergens and allergen-derived peptides in immunotherapy of allergic diseases. *Front Microbiol* 2018; 9:1449.
30. Diesner SC, Bergmayr C, Wang XY, et al. Characterization of *Vibrio cholerae* neuraminidase as an immunomodulator for novel formulation of oral allergy immunotherapy. *Clin Immunol* 2018;192:30–9.
31. Ramirez W, Bourg V, Torralba D, et al. Safety of a proteoliposome from *Neisseria meningitidis* as adjuvant for a house dust mite allergy vaccine. *J Immunotoxicol* 2017;14(1):152–9.

32. Anzaghe M, Schulke S, Scheurer S. Virus-like particles as carrier systems to enhance immunomodulation in allergen immunotherapy. *Curr Allergy Asthma Rep* 2018;18(12):71.
33. Leb VM, Jahn-Schmid B, Kueng HJ, et al. Modulation of allergen-specific T-lymphocyte function by virus-like particles decorated with HLA class II molecules. *J Allergy Clin Immunol* 2009;124(1):121–8.
34. Schmitz N, Dietmeier K, Bauer M, et al. Displaying Fel d1 on virus-like particles prevents reactogenicity despite greatly enhanced immunogenicity: a novel therapy for cat allergy. *J Exp Med* 2009;206(9):1941–55.
35. Focke-Tejkl M, Weber M, Niespodziana K, et al. Development and characterization of a recombinant, hypoallergenic, peptide-based vaccine for grass pollen allergy. *J Allergy Clin Immunol* 2015;135(5): 1207–7.e11.
36. Niederberger V, Neubauer A, Gevaert P, et al. Safety and efficacy of immunotherapy with the recombinant B-cell epitope-based grass pollen vaccine BM32. *J Allergy Clin Immunol* 2018;142(2):497–509.e9.
37. Fanuel S, Tabesh S, Mokhtarian K, et al. Construction of a recombinant B-cell epitope vaccine based on a Der p1-derived hypoallergen: a bioinformatics approach. *Immunotherapy* 2018;10(7):537–53.
38. Senti G, Cramer R, Kuster D, et al. Intralymphatic immunotherapy for cat allergy induces tolerance after only 3 injections. *J Allergy Clin Immunol* 2012;129(5): 1290–6.
39. Zaleska A, Eiwegger T, Soyer O, et al. Immune regulation by intralymphatic immunotherapy with modular allergen translocation MAT vaccine. *Allergy* 2014;69(9): 1162–70.
40. Akdis CA, Blaser K. Bypassing IgE and targeting T cells for specific immunotherapy of allergy. *Trends Immunol* 2001;22:175–8.
41. Rauber MM, Wu HK, Adams B, et al. Birch pollen allergen-specific immunotherapy with glutaraldehyde-modified allergoid induces IL-10 secretion and protective antibody responses. *Allergy* 2019;74(8):1575–9.
42. Marth K, Focke-Tejkl M, Lupinek C, et al. Allergen peptides, recombinant allergens and hypoallergens for allergen-specific immunotherapy. *Curr Treat Options Allergy* 2014;1:91–106.
43. Akdis CA. Therapies for allergic inflammation: refining strategies to induce tolerance. *Nat Med* 2012;18(5):736–49.
44. Prickett SR, Rolland JM, O'Hehir RE. Immunoregulatory T cell epitope peptides: the new frontier in allergy therapy. *Clin Exp Allergy* 2015;45(6):1015–26.
45. Niederberger V, Horak F, Vrtala S, et al. Vaccination with genetically engineered allergens prevents progression of allergic disease. *Proc Natl Acad Sci U S A* 2004;101(Suppl 2):14677–82.
46. Asturias JA, Ibarrola I, Arilla MC, et al. Engineering of major house dust mite allergens Der p 1 and Der p 2 for allergen-specific immunotherapy. *Clin Exp Allergy* 2009;39(7):1088–98.
47. Worm M, Lee HH, Kleine-Tebbe J, et al. Development and preliminary clinical evaluation of a peptide immunotherapy vaccine for cat allergy. *J Allergy Clin Immunol* 2011;127(1):89–97. e114.
48. Spertini F, Perrin Y, Audran R, et al. Safety and immunogenicity of immunotherapy with Bet v 1-derived contiguous overlapping peptides. *J Allergy Clin Immunol* 2014;134(1):239–40.e13.
49. Mosges R, Bachert C, Panzner P, et al. Short course of grass allergen peptides immunotherapy over 3 weeks reduces seasonal symptoms in allergic

- rhinoconjunctivitis with/without asthma: a randomized, multicenter, double-blind, placebo-controlled trial. *Allergy* 2018;73(9):1842–50.
50. Mosges R, Kasche EM, Raskopf E, et al. A randomized, double-blind, placebo-controlled, dose-finding trial with *Lolium perenne* peptide immunotherapy. *Allergy* 2018;73(4):896–904.
 51. Mosges R, Koch AF, Raskopf E, et al. *Lolium perenne* peptide immunotherapy is well tolerated and elicits a protective B-cell response in seasonal allergic rhinitis patients. *Allergy* 2018;73(6):1254–62.
 52. Benito-Villalvilla C, Soria I, Subiza JL, et al. Novel vaccines targeting dendritic cells by coupling allergoids to mannan. *Allergo J Int* 2018;27(8):256–62.
 53. Soria I, Lopez-Relano J, Vinuela M, et al. Oral myeloid cells uptake allergoids coupled to mannan driving Th1/Treg responses upon sublingual delivery in mice. *Allergy* 2018;73(4):875–84.
 54. Hoffmann HJ, Valovirta E, Pfaar O, et al. Novel approaches and perspectives in allergen immunotherapy. *Allergy* 2017;72(7):1022–34.
 55. Curin M, Garmatiuk T, Resch-Marat Y, et al. Similar localization of conformational IgE epitopes on the house dust mite allergens Der p 5 and Der p 21 despite limited IgE cross-reactivity. *Allergy* 2018;73(8):1653–61.
 56. Klunker S, Saggar LR, Seyfert-Margolis V, et al. Combination treatment with omalizumab and rush immunotherapy for ragweed-induced allergic rhinitis: inhibition of IgE-facilitated allergen binding. *J Allergy Clin Immunol* 2007;120(3):688–95.
 57. Larenas-Linnemann D, Wahn U, Kopp M. Use of omalizumab to improve desensitization safety in allergen immunotherapy. *J Allergy Clin Immunol* 2014;133(3):937–937.e2.
 58. Lombardi C, Canonica GW, Passalacqua G. Allergen immunotherapy as add-on to biologic agents. *Curr Opin Allergy Clin Immunol* 2018;18(6):502–8.
 59. Dantzer JA, Wood RA. The use of omalizumab in allergen immunotherapy. *Clin Exp Allergy* 2018;48(3):232–40.
 60. von Moos S, Johansen P, Tay F, et al. Comparing safety of abrasion and tape-stripping as skin preparation in allergen-specific epicutaneous immunotherapy. *J Allergy Clin Immunol* 2014;134(4):965–7.e4.
 61. Kundig TM, Johansen P, Bachmann MF, et al. Intralymphatic immunotherapy: time interval between injections is essential. *J Allergy Clin Immunol* 2014;133(3):930–1.
 62. Devillier P, Dreyfus JF, Demoly P, et al. A meta-analysis of sublingual allergen immunotherapy and pharmacotherapy in pollen-induced seasonal allergic rhinoconjunctivitis. *BMC Med* 2014;12:71.
 63. Matsuoka T, Bernstein DI, Masuyama K, et al. Pooled efficacy and safety data for house dust mite sublingual immunotherapy tablets in adolescents. *Pediatr Allergy Immunol* 2017;28(7):661–7.
 64. Gotoh M, Yonekura S, Imai T, et al. Long-term efficacy and dose-finding trial of japanese cedar pollen sublingual immunotherapy tablet. *J Allergy Clin Immunol Pract* 2019;7(4):1287–97.e8.
 65. Okamoto Y, Fujieda S, Okano M, et al. Efficacy of house dust mite sublingual tablet in the treatment of allergic rhinoconjunctivitis: a randomized trial in a pediatric population. *Pediatr Allergy Immunol* 2019;30(1):66–73.
 66. Masuyama K, Okamoto Y, Okamiya K, et al. Efficacy and safety of SQ house dust mite sublingual immunotherapy-tablet in Japanese children. *Allergy* 2018;73(12):2352–63.
 67. Demoly P, Kleine-Tebbe J, Rehm D. Clinical benefits of treatment with SQ house dust mite sublingual tablet in house dust mite allergic rhinitis. *Allergy* 2017;72(10):1576–8.

68. Okamoto Y, Fujieda S, Okano M, et al. House dust mite sublingual tablet is effective and safe in patients with allergic rhinitis. *Allergy* 2017;72(3):435–43.
69. Ponce M, Schroeder F, Bannert C, et al. Preventive sublingual immunotherapy with House Dust Mite extract modulates epitope diversity in pre-school children. *Allergy* 2019;74(4):780–7.
70. Orengo JM, Radin AR, Kamat V, et al. Treating cat allergy with monoclonal IgG antibodies that bind allergen and prevent IgE engagement. *Nat Commun* 2018; 9(1):1421.
71. Morita H, Tamari M, Fujiwara M, et al. IgE-class-specific immunosuppression in offspring by administration of anti-IgE to pregnant mice. *J Allergy Clin Immunol* 2019;143(3):1261–4.e6.
72. Leuthard DS, Duda A, Freiberger SN, et al. Microcrystalline tyrosine and aluminum as adjuvants in allergen-specific immunotherapy protect from IgE-mediated reactivity in mouse models and act independently of inflammasome and TLR signaling. *J Immunol* 2018;200(9):3151–9.
73. Mao J, Heithoff KA, Koep E, et al. Cost of subcutaneous immunotherapy in a large insured population in the United States. *Curr Med Res Opin* 2019;35(2): 351–8.
74. Shamji MH, Kappen JH, Akdis M, et al. Biomarkers for monitoring clinical efficacy of allergen immunotherapy for allergic rhinoconjunctivitis and allergic asthma: an EAACI Position Paper. *Allergy* 2017;72(8):1156–73.
75. Sugita K, Sokolowska M, Akdis CA. Key points for moving the endotypes field forward. In: Agache I, Hellings P, editors. *Implementing precision medicine in best practices of chronic airway diseases*. London: Elsevier; 2018. p. 107–14.